

Should Antibiotic be used as an Accessory Treatment in Periodontology? A Review

¹Shikha Dhir
² Satendra Sharma
³Shailendra Chauhan
⁴ Aditya Sinha

KD Dental College and Hospital

Abstract:- Treatment of periodontal disease aims at achieving a healthy periodontium and preventing the risk of recurrence / disease progression. Both Surgical and Non-surgical therapy are equally effective in improving periodontal disease. The use of antibiotics in conjunction with periodontal therapy has always been perplexing. This review aims at discussing the rationale behind prescribing the antibiotics with periodontal procedures.

I. INTRODUCTION

Periodontitis has been prevailing in developed as well as developing countries, which affects 20-50% of global population with higher prevalence seen among the adolescents, adults, and older individuals, thus arising as a public health concern. (Nazir, 2017). Periodontal disease has substituted dental caries , being the primary cause of tooth loss. It is strongly associated with a number of other disorders such as low birth weight, heart disease ,and stroke.(Sedlacek and Walker,2007) Periodontal disease is considered to be an infectious disease, with complex microbial aetiology. The existence of virulent microorganisms in the dental plaque biofilm, gives an inflammatory response, which results in tooth loss and destruction of supporting periodontium. (Barça, *et al.*, 2015) Host response involves innate, inflammatory, and

adaptive immune responses. (Silva *et al.*, 2015) Several systemic risk factors related to periodontal disease involve, poor oral habits such as smoking, poor oral hygiene, drug abuse tobacco chewing , systemic factors involving diabetes, puberty , stress maternal infection, preterm birth low birth weight, and preeclampsia. Periodontal disease also increases the risk of cardiovascular disease, and this increase in relative risk reaches to 44% among individuals aged 65 years and over.(Nazir, 2017). Periodontal therapy has seen to improve glycemic control in type 2 diabetic subjects. (Mahuli *et al.*, 2020) Treatment of periodontal disease involves the surgical and non-surgical approaches. (Barça, *et al.*, 2015). Non-surgical treatment involves plaque control with mechanical debridement and periodontal maintenance care, whereas surgical treatment involves resective and regenerative procedures. (Manas D *et al.*, 2009, Graziani *et al.*, 2017) However the mechanical debridement alone, may not be effective in all the cases, thus systemic antibiotics remains the treatment of choice. Considering the periodontal surgical procedures, by their nature carry with them a consequent risk of developing infections, thus adjunctive use of antibiotics gains the attention here. Thus this article focuses on the need of antibiotics to be given with both surgical and non-surgical procedures in periodontology. Fig 1.Use of Antibiotics in adjunction to Non-Surgical therapy (Scaling and Root planing)




Fig1. Antibiotic treatment in Gingivitis and Periodontitis, (Caranza 9th edi, Antibiotic/antimicrobial use in Dental Practice).

Periodontal inflammation occurs as a result of microbes and host microbial interaction. (Manas D *et al.*, 2009) Biofilm formed on tooth surfaces has been implicated as chief culprit in the etiopathogenesis of dental caries and periodontal disease. Continuous and regular disruption of these biofilms is essential for prevention and management of periodontal diseases. (Chandki *et al.*, 2011) Scaling and root planing will always remain a gold standard treatment for the disruption of supragingival as well as subgingival biofilm and calculus. (Feres *et al.*, 2015) However SRP is not always successful in reducing the subgingival microbial load completely due to the presence of deep and narrow pockets, due the irregular course of pocket in an individual teeth leading to an irregular base of furcation mainly in multirrooted tooth , grooves and depression onto root surface, or due to "line angles", i.e., the area where a change would be made from one curette blade to another during the root planing process.(Pluss EM, *et al.*,1992)The results of various studies, being conducted for the evaluation of clinical effects of SRP suggest that SRP doesn't show major improvement in all subjects, especially in cases with severe periodontal disease which included deep periodontal pockets and in patients with systemic disease . (Feres *et al.*, 2015) This is probably because of the persistent infections caused by *P.gingivalis*, *A.actinomyetemcomitans* and *P.intermedia*, which have the tendency to invade the soft tissue wall of periodontal pocket and are difficult to eliminate with SRP.(John Walters and Pin-Chuang Lai, 2015) In such case, antibiotics may be required to control the microbial load causing infections along with mechanical debridement as to further 'eliminate or markedly suppress the specific microorganisms which can cause breakdown of periodontal attachment in susceptible patients' (fig 2). (Mombelli, 2017) Hererre *et al.*, at the fourth European Workshop, in a systemic review elaborated that systemic antibiotics can offer an additional benefit when used in conjunction with scaling and root planing, similar results were presented by Haffajee *et al.* (Heitz-Mayfield, 2009) Thus the inability of SRP in removal of microbial aetiology of the infectious periodontal disease, provides the rationale for the use of antibiotics as an adjunctive to periodontal therapy. (Barça, *et al.*, 2015)

Systemic periodontal antibiotic therapy along with mechanical periodontal treatment aims to support the host defense system in overcoming the infection by killing subgingival pathogens that get left after conventional mechanical periodontal therapy. Result of several studies

describe that SRP along with proper oral hygiene measures ,are efficient in preventing periodontal breakdown in most of the individuals by suppressing total supra-subgingival microbes. (Graziani *et al.*, 2017, Wolf, 2004) However, despite smart dental therapy, some individuals continue to experience periodontal breakdown, as major periodontal pathogens like *Porphyromonas gingivalis*, *Aggregatibacter actinomyetemcomitans*, *Fusobacterium-nucleatum*, *Treponemadenticola*, *bacteroids*, intent to invade periodontal tissues or reside in deep pockets or root furcation defects or root concavities or grooves, the sites which are difficult to reach with periodontal instruments, or due to poor host defense mechanisms.(Wolf ,2004, Guzeldemir-Akcakanat,2019)





Fig. 2. Diagrammatic represents the clinical significance of microbial colonization in pocket for initial cause-related periodontal therapy (left), immediately thereafter (middle) and during maintenance (right). ASA, adjunctive systemic antibiotics; BR, biofilm removal; OH, oral hygiene to interfere with recolonization; LAT: local antimicrobial treatment; SRP, scaling and root planing to remove calculus (purple) and biofilm (red).(Mombelli, 2017)

Antibiotics used to treat periodontal disease (listed in table 1), are selected based on the patient's medical and dental status, current medications, and results of microbial analysis, if performed. The prime candidates for systemic antimicrobial therapy are those patients exhibiting attachment loss or having residual pockets even after adequate conventional therapy, or patients with active periodontal disease or associated with predisposing medical conditions. (Graziani *et al.*, 2017, Kapoor *et al.*, 2012, Slots, 2004)

Table 1: Common Systemic Antibiotics in Periodontal therapy

Systemic Antibiotics	Route	Primary Excretion	Doses and Duration
Penicillin group			
Ampicillin	Oral	Renal	250mg-500mg/6h/5-7 days
Amoxicillin	Oral	Renal	500mg/8h/8 days
Penicillinase – resistant penicillin Group			
Cloxacillin	Oral	Renal	500mg/6h/
Dicloxacillin	Oral	Renal	500mg/6h/
Cephalosporins			
Cephalexin	Oral	Renal	250-500mg/6h/5-7days
Tetracycline group			
Tetracycline HCL	Oral	1°Renal	250mg/6h/2-7 years

		2°Biliary	
Minocycline	Oral	1°Renal 2°Biliary	200mg 1 st day then, 100mg/12h/ 21 days
Doxycycline	Oral	1°Renal 2°Biliary	200mg 1 st day then, 100mg/12h/21 days
Erythromycin group			
Clindamycin	Oral	Hepatic	300mg 1 st day, then 150mg/8h/8days
Metronidazole	Oral	Renal	500mg/8h/8days
Azithromycin	Oral	Hepatic	500 mg/qid./3-7 days



Combination therapy of antibiotics (synergism)	Chart 1 Dosage
Metronidazole + amoxicillin	250mg/t.i.d./8 days of each drug
Metronidazole+ ciprofloxacin	500mg/b.i.d./8 days of each drug

Patients with acute periodontal infections (periodontal abscess, acute necrotizing gingivitis/periodontitis and acute necrotising somatitis) may also require antibiotic therapy. However patients having gingivitis or mild-moderate periodontitis usually respond well to mechanical debridement and need not require additional benefit from systemic antibiotic therapy, if required topical antibiotics can be used (Slots, 2004) Moreover microbial samples can be obtained from individual active pockets or from subgingival sites, which will provide a better representation for the range of periodontal pathogens to be targeted with antibiotic therapy. (Kapoor *et al.*, 2012). However systemic antimicrobials are more effective when the biofilm has already been reduced with proper debridement. which should be carried out in the shortest possible time span (approximately 1 week) to achieve effective drug serum levels on the day of debridement completion. Thus mechanical biofilm disruption must always be combined with proper oral hygiene measures performed by patient. (Barca *et al.*, 2015)




Chart 3– Acute gingival infections requiring Antibiotics

Antibiotics as adjunctive to periodontal surgery

When periodontal surgery is performed, post-surgery is often at risk of developing complications including infection, pain, reduced mouth opening, reduced masticatory capability, swelling of variable degree and delayed wound healing. (H.R et al., 2006, Rajpal et al., 2015) However the prevention of infections after the periodontal surgery, by prescribing antibiotics have always been controversial. (H.R et al., 2006) Various meta-analysis studies,



Chart 4:- show that the used of systemic antimicrobials did not show any statistically significant results in relation to the improvement in periodontal parameters. However many studies suggest that antibiotics prescribed postoperatively are effective in reducing the postoperative inflammation, infection and improving wound healing . (H.R et al., 2006, Trivedi, 2014)

As postoperative infection can have a significant effect on the surgical outcome, preventive measures like strict aseptic protocol, anti-infective measures like proper sterilization, disinfection, barrier techniques need to be taken. If such measures are taken, low rate of postoperative infection following periodontal surgery can be seen, thereby avoiding the need for using antibiotics as a postoperative care. (Monica and Biir, 2019) But in an Indian setting attaining an aseptic environment for a clinical study, in every patient is a goal we dream to achieve. (Trivedi, 2014) Thus antibiotic prophylaxis are to be prescribed post operatively mostly in patients who present with a high risk of infection or when traumatic surgical procedures have been performed. (Rajpal et al., 2015).

II. CONCLUSION

Periodontal therapy involves both nonsurgical and surgical to treat periodontal disease. However, no periodontal treatment has shown clear superiority over any other periodontal treatment. Treatment of periodontitis involves a fine balance of skilled techniques and a better access to the treatment site, which together decrease the risk of disease progression. (Manas *et al*, 2009, Graziani *et al.*, 2017) In particular instrumentation, typically performed using root debridement with sonic/ultrasonic instruments and the finishing phases of root planing with manual curettes may not always be effective in complete removal of subgingival bacteria. Thus antibiotics can be used as an adjunctive, in the patients who do not respond to mechanical treatment, patients with acute or severe periodontal infection, and patients who are systematically compromised. (Barça, *et al.*, 2015, Graziani *et al.*, 2017). Additionally systemic antibiotics may be associated with significant adverse effects and increased bacterial resistance, thus should be prescribed only for indicated patients and with caution in patients on long-term medication for cardiovascular disease, asthma, seizures, or

diabetes, as there can be drug interactions. (Barça, *et al.*, 2015), Graziani *et al.*, 2017] As antibiotic resistance constitutes an increasing problem, anti-microbial susceptibility testing of isolated pathogens is important, if unavailable, combination therapy is preferred against the mixture of pathogenic bacteria, in the periodontal lesions. (Barça, *et al.*, 2015) Although local delivery of antimicrobials into a periodontal pocket can also improve the periodontal health. However these drug systems do not provide a superior result when compared to scaling and root planing and should be used as an adjunctive to SRP. (Rajpoot AS *et al.*, 2017) .However, prophylactic antibiotic therapy, as recommended by the American heart Association, is mandatory for patients with specific medical problems such as immunodeficiency, metabolic diseases, irradiated in the head and neck area, those at high and moderate risk for endocarditis and for extensive or prolonged surgery

REFERENCES

- [1]. Amitha R, Agumbe P, Biju T (2016). Local Drug Delivery in periodontal diseases. A Review. *NUJHS* 6: 74-79.
- [2]. Anoop K, Rajan M, Vishaka G, Deepak G (2012). Systemic antibiotic therapy in periodontics. *Dent Res J (Isfahan)* 9: 505–515.
- [3]. Ankur SR, Anuj SP, Swapnil T, Kapil C, Preeti R, Ambika C(2017). Local drug delivery in periodontics. *IJRHAS* 3:63-67
- [4]. Andrea M (2017). Microbial colonization of the periodontal pocket and its significance for periodontal therapy. *Periodontol* 2000 0; 1-12.
- [5]. Arab H, Sargolzaie N, Moeintaghavi A Ghanbari H, AbdollahiZ (2006). Antibiotics to Prevent Complications Following Periodontal Surgery. *IJP* 2:205-208.

- [6]. Catherine P, Fareeha B, Isaac M, Pascale S, Nadia B, Olivier H(2019). Contribution of Statins towards Periodontal Treatment: A Review. *MEDIATINFLAMM* 2-33.
- [7]. Darshan B, Suchetha A, Karthik V, Akanksha G (2011). Newer trends in Local Drug Delivery for Periodontal Problems –A Preview. *IJCD* 2:59-62.
- [8]. Deepa S, Pavithra. D (2020). Local Drug Delivery in Periodontics - A Review. *EJMCM* 7:1874-1881.
- [9]. Dieter D (2017). The periodontal pocket: pathogenesis, histopathology and consequences. *Periodontol* 2000 0:1-8.
- [10]. Diaz PI, Kolenbrander PE (2009).Subgingival Biofilm Communities in Health and Disease. *Rev. Clin. Periodoncia Implantol. Rehabil. Oral* 2:187-192.
- [11]. Ece B, Emine C, Serdar C (2015). Adjunctive use of antibiotics in periodontal therapy. *Istanbul Univ Fac Dent* 49:55-62.
- [12]. Filippo G, Dimitra K, Bettina A, David H (2017). Nonsurgical and surgical treatment of periodontitis: How many options for one disease? *Periodontol* 2000 75: 152–188.
- [13]. Guzeldemir.A (2019).Systemic Antibiotics in the Treatment of Periodontitis. *Dent Med Res* 7:33-4.
- [14]. Herbert FW, Thomas MH (2006). Color atlas of dental hygiene-periodontology. *Microbiology—Biofilm* 24-29.
- [15]. Jaisika R, Aakash A, Ruchika P, Madhav MG,(2015).Preventing post-operative swelling after periodontal surgery. *J Oral Res Rev* 7:31-4.
- [16]. Jack GC , Gary A , Tord B, Iain LC, Soren J, Kenneth S *et al.*(2018).A new classification scheme for periodontal and peri-implant diseases and conditions – Introduction and key changes from the 1999 classification. *Clin Periodontol.* 45 :(Suppl 20) S1–S8.
- [17]. Jingou L, Xinyu P, Xuedong Z, Jing Z, Lei C (2020). Emerging Applications of Drug Delivery Systems in Oral Infectious Diseases Prevention and Treatment .*Molecules* 25:2-29.
- [18]. John W, Chuang L (2015) Should Antibiotics Be Prescribed to Treat Chronic Periodontitis? *Dent Clin N Am* 2-15.
- [19]. Lja HM (2009). Systemic antibiotics in periodontal therapy. *Aust Dent J* 54:96–101.
- [20]. Monica GS, Biir M.S.M (2019).Evaluation of the Role of Antibiotics Following Periodontal Surgery in Chronic Periodontitis Patients in Preventing Post-Operative Complications-A Relative Clinical Study. *Int.Res.J.Med.Sci*1; 71-76.
- [21]. Magda F, Luciene C, Geisla M & Marcelo F, (2015).Systemic antibiotics in the treatment of periodontitis . *Periodontol* 2000 67:131-186.
- [22]. Marie B, Andrew H, Charles HJ, Blaine AP (2018). Phenotypic Variation during Biofilm Formation: Implications for Anti-Biofilm Therapeutic Design. *Materials* 11:2-18
- [23]. Manas D, Srinivas SR, Jithendra KD (2009). Role of antibiotics in periodontal therapy. *BJMS* 8:91-101.
- [24]. Meena R, Angela G, Kevin D (2007).Locally Delivered Antimicrobials In The –Management Of Periodontitis: A Critical Review Of The Evidence For Their Use In Practice *DENT UPDATE* 34: 494-506.
- [25]. Muhammad AN (2017). Prevalence of periodontal disease, its association with systemic diseases and prevention. *IJHS* 1:2.
- [26]. Nora S, Loreto A, Denisse B, Nicolás D, Jocelyn GS, Rolando V, Marcela H, *et al.* (2015). Host response mechanisms in periodontal diseases. *J Appl Oral Sci* 23:329-55.
- [27]. Paul E, Roxanna N , David S, Paul G , Jamie S., Robert J (2002).Communication among Oral Bacteria *Microbiol. Mol. Biol. Rev.* 66:486-505.
- [28]. Paul E, Jack L (1993).Adhere Today, Here Tomorrow: Oral Bacterial Adherence. *J.Bacteriol* 175 : 3247-3252.
- [29]. Pragati S, Ashok S, Kuldeep S (2009).Recent advances in periodontal drug delivery systems. *IJDDT* 1:1-14.
- [30]. Rajeshwari H, Dinesh D, Satveer J , Meghana R , Kiran J ,Shabana S *et al.*(2019). Local drug delivery systems in the management of periodontitis: A scientific review. *JCR* 307:393-409.
- [31]. Rateitschak EM, Schwarz JP, Guggenheim R, Duggelin M, Rateitschak KM (1992) .Non-surgical periodontal treatment: where are the limits? An SEM study. *J Clin Periodontol* 19: 240-244.
- [32]. Rita C, Priyank B, Ruchi B (2011).Biofilms: A microbial home. *J Indian Soc Periodontol* 15:111-114.
- [33]. Sahilhusen I, Mukesh R (2014). Pharmaceutical Controlled Release Drug Delivery Systems: A Patent Overview. *J Drug Design Pharmacol* 1:2-22.
- [34]. Sedlacek, MJ, Walker C (2009), Antibiotic resistance in an *in vitro* subgingival biofilm model. *Oral Microbiol Immunol* 22: 333–339.
- [35]. Shilpa T (2008). Antibiotics and periodontal surgery. *Clin Periodontol* 5:58-63.
- [36]. Simpy AM, Ali MZ, Mohammed AJ, Alnomari S, Gargi S, Hosam AB, *et al.* (2020). Antibiotics for Periodontal Infections: Biological and Clinical Perspectives. *J Contemp Dent Pract* 21:372–376.
- [37]. Slots J (2004).Systemic Antibiotics in Periodontics. *J Periodontol* 75:1553-1565.
- [38]. Thomaes R, Slots J (1996). Local delivery of antimicrobial agents in the periodontal pocket. *Periodontol* 2000 10:139-159.
- [39]. Yu H, Xiaoyu H, Xuedong Z , Mingyun Li , Biao R, Xian P *et al.*(2018). Influence of Dental Prosthesis and Restorative Materials Interface on Oral Biofilms. *Int. J. Mol. Sci.* 19:2-17