

Valuation of Awareness and Precautionary Practice of Corona Virus among Communities; the Case of Kaffa Zone, South-Western Ethiopia

Getachew Roba Agegneu^{*1}; Seyoum Robo G/Mariam¹; Abera Haile Adello¹; Yasin Abafita Abawari¹; Abiyot Mamo Mergia¹ and; Habtamu G/Senbet Sahile¹

¹⁾ Department of Psychology; Bonga University, Bonga, kaffa, 334, Ethiopia

¹⁾ Department of Biology; Bonga University, Bonga, Kaffa, 334, Ethiopia

¹⁾ Department of Chemistry; Bonga University, Bonga, Kaffa, 334, Ethiopia

¹⁾ Department of Sport Science; Bonga University, Bonga, kaffa, 334, Ethiopia

¹⁾ Department of mathematics; Bonga University, Bonga, Kaffa, 334, Ethiopia

¹⁾ Department of Biology; Bonga University, Bonga, kaffa, 334, Ethiopia

Abstract:- The severe acute respiratory syndrome coronavirus and resultant coronavirus disease (COVID-19) have evolved into a pandemic, requiring persons around the world to attend to rapidly changing messages about public health and take immediate actions. Related to this, community's awareness toward the preventive practice of coronavirus is assessed. Hence, the purpose of this was to investigate the awareness and preventive practice among communities. Cross-sectional survey design was conducted among 400 participants. Simple random sampling, technique particularly lottery method was used to select the study participants. Data was collected by using questionnaire and observation. After the data was collected, data entry and analysis were done using SPSS version 26. Hence, data gathered using questionnaire and observation was computed by using simple descriptive statistics (percentage and frequencies). Regarding awareness level of communities 61% of respondents aware that covid-19 transmits from victim by hand shaking and close contacts. And hence, 37.8% of participants aware that awareness creation was done by health organization service centers. And hence, the findings in this study revealed that community's awareness was high on each specific aspect of COVID-19 among the study participants. Communities are not still totally pay attention to aware about COVID 19, crisis. According to finding in this research communities wash their hand in order to prevent covid-19 better than the previous. Kaffa zone as well as stakeholders should have to provide ample information for all communities including urban and rural societies to aware about COVID-19 and practice how to prevent.

Keywords:- Awareness; Communities; Corona Virus; Precautionary and Valuation

I. INTRODUCTION

In earlier December, first case of pneumonia of unknown cause originated in Wuhan, capital city of Province Hubei, China, and on 31 December 2019, with emergence of more such cases, Wuhan gained attention by World Health Organization. (Huang C, 2019).

The pathogen identified was named as novel coronavirus (2019-nCoV), currently called as severe acute respiratory syndrome corona virus-2 (SARS-CoV-2), an enveloped and single stranded RNA virus (Guan W (2019) which has phylogenetic resemblance to SARS-COV-1. Lu R *et al.*, (2019) Owing to rapid spread of this deadly virus from epicenter to number of countries, WHO declared it as public health emergency of international concern (PHEIC) on January 30, 2020. Later, due to uncased fast spread, severity of illness, the continual escalation in number of affected countries, cases and causalities, WHO declared coronavirus disease 2019 (COVID-19) a global pandemic on 11 March 2020 WHO, (2020). To date (12 April, 2020), the COVID-19 have spread to 210 countries and territories accounted for 1,790,550 laboratory confirmed cases and 109,654 mortalities also attributed to this deadly pathogen Worldometer 2020:1–22.

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and resultant coronavirus disease 2019 (COVID-19) have evolved into a pandemic, requiring persons around the world to attend to rapidly changing messages about public health and take immediate actions to minimize their risk for infection and the spread of the virus (Worldometer, COVID-19: 2020). This unprecedented global crisis has also been marked by miscommunication regarding the imminent threat of COVID-19, leading to public confusion and inaction (Ioannidis, J. P. A., (2019).

COVID-19 transmits from person to person by droplets when an infected person sneezes and by direct contact and the virus has an incubation period of 4-14 days (CDC: (2020). Elderly and patients who suffered with chronic medical conditions like diabetes and cardiovascular diseases are more likely to get severe infection. The main manifestations of COVID-19 are fever, dry cough, dyspnea, myalgia, fatigue, hypolymphemia, and radiographic evidence of pneumonia. Complications (e.g., acute respiratory distress syndrome (ARDS), arrhythmia, shock, acute cardiac injury, secondary infection, and acute kidney injury) and death may occur in severe cases Lei S et al. (2020). Presently, no antiviral therapy or vaccine is explicitly recommended for COVID-19 and implementation of preventive measures to control COVID-19 is the mainstay critical intervention. Bhagavathula AS, Aldhalei WA, Rahmani J, Mahabadi MA, Bandari DK. (2020).

The COVID-19 symptoms can range from mild (or no symptoms) to severe illness and are mainly characterized by fever, dry cough, dyspnea, headache, sore throat and rhinorrhea and sometimes hemoptysis (Adhikari *et al.*, 2020, Cascella, Rajnik *et al.*, 2020). The main route of transmissions is close contact (about 6 feet or two arm lengths) with a person who has COVID-19, respiratory droplets when an infected person coughs, sneezes, or talks and touching a surface or object that has the virus on it, and then by touching mouth, nose, or eyes (Guo *et al.*, 2020).

The same factors that make individuals more vulnerable are also associated with reduced ability to access and understand health information, make well-informed decisions, and take optimal health-promoting actions a skill set commonly called “health literacy” (Wolf MS, Gazmararian JA, Baker DW 2005; Paasche-Orlow MK, Parker RM, Gazmararian JA, *et al.*, 2005). This is especially true when the health information itself is not timely, trusted, consistent, or actionable. Health literacy has emerged over the past 3 decades as one of the strongest psychosocial determinants of health, and it has also been shown to explain a range of health disparities by age, race/ ethnicity, and socioeconomic status (Mantwill S, Monestel-Uman˜a S, Schulz PJ, 2015). Thus, in uncertain times like this, when the interpretation of critical and ever-changing public health messages is paramount, many vulnerable populations may be further marginalized by inadequate health communication, posing substantial risks to themselves and their communities. Literature suggests that lack of knowledge and misunderstandings among People and professionals leads to delayed diagnosis, spread of disease and poor infection control practice (Saqlain M., 2020).

The novel coronavirus (CoV) was initially identified in Wuhan, China, in December 2019 among a group of patients that presented with an unidentified form of viral pneumonia with common history of visiting a seafood market. (Zhu N. *et al.*, 2020)

Ethiopia reported its first case of COVID-19 on March 13, 2020. (First Case of COVID-19 Confirmed in Ethiopia.). Since then, gradual increment in the number of cases has been reported by the ministry of health. In countries like Ethiopia, where low surveillance and laboratory capacity, scarcity of healthcare human resources, and inadequate financial capacity are defining features of the healthcare system, the outbreak of a cureless viral infection with no vaccine will have a devastating impact. (Nkengasong JN, Mankoula W. Looming, 2020). Due to this, limiting the rate and the extent of COVID-19 outbreak is the best strategy to minimize public health and socioeconomic impact of the pandemic in the country, and healthcare professionals play the central role. (Voo T, Capps B. 2010).

Many efforts have been made by the government of the country and the ministry of health in order to prevent the spread of the disease. Contagious transmission route of the disease made the Universities and schools to be closed; private worker and government employees were assigned to stay at home; transportation was stopped for shorter period and re allowed to work with 50% capacity. Marketing areas were redistributed; Universities and large hall, including Millennium halls were assigned to get ready COVID-19 patient treatment. All social media both private and government played an unquantifiable role to creating awareness. Kaffa zone is one of the zones found in southern regional state, indicates that there is opportunity for spreading of the disease.

To facilitate an urgent response to potential COVID-19 outbreak in Ethiopia, it is important to evaluate the awareness and preventive practice of health care workers who engage in the response against the outbreak. This study was launched to assess COVID-19-related awareness preventive practice among Kaffa zone south western Ethiopia. Hence, there are many chances such as social interaction among the people; daily transport which moves from central part of country, Addis Ababa to the area and it is also surrounded by other neighboring regions and zone. On the other hand, the extent of awareness, attitudes and effectiveness of preventive practice toward the disease, is not known. Many researches had been conducted on the novel Covid-19 by encompassing different variables; for example, Zelalem T, Malede, Y, Zenebe N and Akeberegn A, 2020 and Muhammad S *et al.*, 2020 conducted research merely on the health professionals, but our research emphasizes on the community; beyond to this; the researchers collected data from different woredas where different social issues were taken such as ceremonies (funeral and wedding) were considered.

- How much of the member of the community do have awareness on coronavirus disease?
- What the awareness of community members aware coronavirus disease?
- What is the extent of corona virus preventive practices that were utilized in the study area?

II. MATERIALS AND METHODS

The study was carried out in Kaffa zone, southern nations, nationalities and people regional state, southwestern Ethiopia. The capital town of the zone, Bonga, is located 449 km southwest of Addis Ababa, capital city of Ethiopia. The name of the study area “Kaffa” is well known and is every day on the minds of the many people throughout the globe who savor the juice extracted from the berry of a plant which originally grew in Kaffa, coffee (Grühl, 1935). Kaffa zone is bordered by Oromiya region on the north, Semen Omo zone on the east, Debub Omo zone in south east, Bench Maji zone on south and south west and Shaka zone on the west. The zone is subdivided into 12 districts, namely Chena, Bitta, Gesha, Decha, Gimbo, Goba, Cheta, Gawata, Menjiyo, Saylem, Shishinda, Tello, and two administrations.

The Constraints that the researchers had faced during the time of conducting the study were so serious and be challenges to conduct research. The most limitation was that majority of participants need to hand shake and embrace the data collector, some other participants ask for financial support to give real information about what is asked, some participants who are aware about the transmission of covid-19 had denied to take the questionnaire and give the data because of the fear of contact with the other people because they have been aware that covid-19 transmits with contact, certain participants are not providing the answer for the question listed on the questionnaire i.e. there are incomplete answers etc.4

The study design for this study was cross-sectional survey design. This is preferred because the means that the

researcher use to obtain data from participants was questionnaire and observation that was administered at one time to investigate the awareness and preventive practice of corona virus (covid-19). among communities; the case of Kaffa zone, south western Ethiopia.

Yamane (1967) was used to calculate sample sizes, n= 400. This formula was used to calculate the sample sizes at 95% confidence level and, e is the desired level of precision 0.05, total population

$$N = 874,716 \text{ based on 2007 CSA data}$$

$$n = \frac{N}{1 + N(e)^2}$$

The study population eligible for participation in this survey were two city administrations, (Bonga and Chena town) for their high traffic movements and three woredas Gimbo, Saylem and Addiyo for their boarder due there may be entry of patient or the disease. After selection of woredas two kebeles from each woreda and town administration had participated about 40 respondents were eligible by lottery method.

After the data was collected, data entry and analysis were done using SPSS version 26. Hence, data gathered using questionnaire and observation was computed by using simple descriptive statistics (percentage and frequencies).

III. RESULT

Table 1 Socio- Demographic Characteristics of the Respondents, N=400

Variables	Categories	Male	Female	Total	Percent (%)
Sex	Male/Female	296	104	400	100
Age	< 30	75	31	106	27
	31-45	161	49	210	52.5
	46-60	49	20	69	17.25
	60+	11	4	15	3.75
Educational status	1-8	35	19	54	13.5
	9-12	52	29	81	20.25
	12+	167	39	206	51.5
	Illiterate	42	17	59	14.75
Work	Private	94	48	142	35.5
	Gov.t employee	202	56	258	64.5
Religion	Orthodox	123	28	151	37.75
	Muslim	54	22	76	19
	Catholic	20	15	35	8.75
	Protestant	88	31	119	29.75
	Others	11	8	19	4.75
Marital Status	Married	186	68	254	63.5
	Single	110	36	146	36.5

A total of four hundred individuals were participated in this study. The objective of this study was to investigating the awareness and preventive practice of community. Hence, the total number of the distributed questionnaires were 400 and all questionnaires were filled completely and consistently with a response rate of 100%. Among the total respondents who filled the questionnaire 296 (74%) were males and the

rest 104 (26%) were females. The majority of respondents' age was on the intervals of 31-45. As observed from above table majority of respondents 206 (51.5%) were above grade 12th, hence, 258 (64.5%) of participants were government employee, 151 (37.75%) of respondents were orthodox religion followers and finally 254 (63.5%) of the respondents were married in their marital status.

Table 2. Awareness level about the Corona virus (COVID-19)

Item	Frequency	Percentage (%)
It transmits from victim by hand shaking and closes contact	244	61.0
It caused by eating uncooked foods	33	8.3
It Transmit by sneezing from victim to health person	92	23.0
No clue for its transmission	31	7.8
Total	400	100

As it is displayed from the above table it assesses the awareness of the community in the study area and hence, 244(61%) of the participants replied that covid-19 transmits from victim to other by hand shaking and closet contact. Related to this the community have awareness about the transmission of COVID-19. Related to this 33 (8.3%) of the participants responded that coronavirus transmits by eating

uncooked foods, whereas 92(23%) of the respondents replied that covid-19 transmits by sneezing from victim to healthy person. However, 31(7.8%) of respondents replied that they did not have hint about the transmission of covid-19. Generally, in this context we understood that majority of the community aware that covid-19 transmits by hand-shaking and close contact.

Table 3. Group of individuals affected by Corona virus

Item	Frequency	Percent (%)
Upper age groups	57	14.2
Affects age independently	321	80.3
I have no awareness	22	5.5
Total	400	100

The above table demonstrates the age group that covid-19 affects. Hence, 57 (14.2%) of respondents aware that a covid-19 pandemic disease affects the upper age groups, 321(80.3%) of respondents replied that they have awareness that covid-19 affects age independently, while 22(5.5%) of

respondents replied that as they have no awareness that which age group covid-19 affects. Here in this case, we understood that the age group affected by covid-19 were the upper age group; this in other way does not mean that the other age group were not affected by covid-19.

Table 4. Extent of awareness level toward covid-19

Variable	Frequency	Percent (%)
Sufficient	207	51.7
Not sufficient	193	48.25
Total	400	100

The above table displayed that the community's awareness level toward covid-19. Related to this 207 (51.7%) of participants replied that their awareness level is sufficient and they have good awareness about covid-19. While 193(48.25%) of respondents replied that their awareness is

not sufficient and they did not have much enough awareness about covid-19. Hence, the community's awareness level is sufficient about the COVID-19 in general. However, the practical implementation needs care.

Table 5. Awareness creation

Response	Frequency	Percentage (%)
Yes	293	73.3
No	107	26.8
Total	400	100

As it illustrated from the existence of community awareness creation; related to this 293 (73.3%) of the respondents responded that as there is awareness creation in their community while 107 (26.8%) of participants replied

that there is no awareness creation program in their community. In this case community's awareness creation was made by Kaffa zone government medias.

Table 6. ways by which community awareness was created

Ways	Frequency	Percent (%)
By Radio	138	34.5
In Health organization service	151	37.8
In Idir/market	108	27
Total	400	100

As demonstrated from the above table 6 the ways of community awareness creation were assessed. Related to this 138 (34.5%) of participants replied that the awareness about covid-19 was created through radio, while 151 (37.8%) of participants replied that their awareness toward covid-19 was

created through health organization service centers and 108 (27%) of participants replied that their awareness toward covid-19 was created when they are involving Idir as well as market places.

Table 7. Belief that keeping physical distancing is effective

Places	Frequency	Percent
In shops	57	14.2
In market	44	11
In lamentation	11	2.8
All	288	72
Total	400	100

As it is displayed from the above table, the community's belief of keeping physical distancing is effective; related to this 57 (14.2%) of respondents replied that they have confidence keeping physical distancing is effective in shops, while 44 (11%) of respondents believe and replied that keeping physical distancing in market places is effective, 11 (2.8%) of participants replied that keeping physical

distancing in crying is effective not to be affected by COVID-19. And lastly, 288(72%) of respondents replied that all the above places have high opportunity to transmit the disease. From the above table, it is real that physical distancing is effective in different aspects of the community's day to day activity.

Table 8. Duration of awareness creation provided

Periods	Frequency	Percent (%)
Many Times,	147	36.8
Twice	53	13.3
Rarely	163	40.8
Once	36	9
Total	400	100

The above table demonstrates the duration of awareness creation provided in the community; beside to this 147(36.8%) of the participants responded that awareness creation is provided many times, 53(13.3%) of participants responded that awareness creation is provided twice a week, 163 (40.8%) of participants replied that awareness creation

was provided rarely and finally 36 (9%) of respondents responded that in their community awareness creation is provided once a week. The other issue is duration of awareness creation that majority of community aware that awareness creation was made many times by different medias in different dimensions of the communities.

Table 9. Corona virus disease can be protected only with belief alone without any suggested protection ways

Variable	Frequency	Percent
strongly disagree	201	50.2
disagree	71	17.8
undecided	62	15.5
agree	44	11.0
strongly agree	21	5.3
Total	400	100.0

As it is displayed by the above table, participants awareness assessed and with respect to this 201 (50.2%) and 71 (17.8%) of participants replied that they strongly disagree and disagree respectively that Corona virus disease cannot be protected only with belief alone without any suggested protection ways. Contrary to this 44 (%) and 21 (5.3%) of

respondents replied that they agree and strongly agree with the point Corona virus disease can be protected only with belief alone without any suggested protection ways. Here from the above case participant believe that covid-19 may not be protected solely by belief without any determined protection ways.




Figure 1. Prevention methods used by families

Among different methods used for prevention of covid-19 by local people in the study area more than 50%’s use water and soup, whereas face mask users are fewer in number.




Figure 2. Do you use soup and water in your outdoor?

As shown in chart above about 188(47%) of the respondents do not practiced using soup and water as a preventive measure; 184(46%) rarely use and the rest only about 28(7%) of the population always use prepare water and soup out door in their home area.

Table 17. Staying at home and avoid hand shaking cannot protect Corona virus disease

Item	Frequency	Percent
Strongly disagree	176	44.0
dis agree	25	6.3
Undecided	81	20.3
Agree	50	12.5
strongly agree	66	16.5
Total	400	100

In prevention of staying at home issue and avoiding at home to prevent covid-19; majority or 176(44%) of the respondent strongly disagree, and the list percent's 25(6.3%) disagree; 81(20.3%) respondents undecided; 50(12.5%) agreed; 66(16%) are strongly agreed on the idea of staying at home and handshakes cannot prevent corona virus disease. Finally, majority of the communities agree that by staying at home they can prevent corona virus transmission and overall effect on their health situations.

IV. DISCUSSION

As of August 23, 213, 489 confirmed cases of COVID-19 had been reported worldwide, including 804,556 deaths. In Saudi Arabia, 104,000 cases have been confirmed. In our study, most participants had a high level of knowledge regarding COVID-19. Indeed, 61% of participants aware that COVID-19 transmits from victim by hand shaking and closes contact and 80.3% of participants aware that COVID-19 age independently respectively. This high level of knowledge was resulted from the Ethiopia's MOH's efforts to increase public knowledge of this novel virus by sending mobile messages and using different means of communication, including television and social media. A consistent study that Zhong et al conducted on Chinese residents also assessed participants' knowledge of COVID-19. (Zhong B, Luo W, Li H, et al.,2020). During the rapid rising era of the COVID-19 outbreak, the majority of communities had good information, optimistic attitudes, and suitable procedures about COVID-19. Furthermore, good COVID-19 knowledge is linked to positive attitudes and safe COVID-19 practices, implying that health education programs aiming at enhancing COVID-19 knowledge can assist encourage positive attitudes and safe COVID-19 practices.

Hence, majority of residents in this study findings suggest that Chinese residents of a relatively high level of socioeconomic status, in particular women, have had good knowledge, optimistic attitudes, and appropriate practices towards COVID-19 during the rapid rise period of the COVID-19 outbreak. Particularly their knowledge of the clinical characteristics and prevention of the virus. In their study, the authors reported that most of the respondents (90%) were knowledgeable about COVID-19, consistent with the results of our study. Hence, in order to develop the awareness of communities about COVID-19 and its preventive practice in their day-to-day life activity several

activities are done; such as announcement about prevention of COVID-19 with different social media, television, radio, presentations about COVID-19 with microphones where peoples are accumulated in one area such as market place, educational setting, funerals, weeding, night parties etc. Similarly, a study by Peng et al that assessed Chinese University students' knowledge regarding COVID-19 reported that most of the participants (82.34%) had a basic knowledge of this disease (Peng Y, Chenchen P, Zheng Y, *et al.*, 2020). In other study conducted in China assessed the Chinese knowledge about coronavirus; regarding with this China community have good knowledge about the virus. Hence, Chinese government had played a great role in developing the community's awareness by different medias. Aside from that, the government plays an important role in raising community awareness and attitudes by disseminating important information through mechanisms such as television, social media, barouches, and magazines, among others, to raise community awareness and attitudes about how to avoid being afflicted by the virus. Another study by Hussain et al reported that Nepalese residents exhibited satisfactory knowledge (Hussain A, *et al.*, 2020). A similar study that Abdelhafiz et al assessed Egyptians' overall awareness' toward and knowledge and perceptions of this novel coronavirus disease and reported that the participants generally had good knowledge, which is consistent with our study; therefore, according to Annune, A.E *et al.*, 2020) study, when an individual comes in contact with an infected person, there are high chances of an individual becoming a victim and the end result in most cases is death as no vaccine has being discovered yet. But not only direct and closed contact with victim spread this disease, but also contacting anything that the victim contacted with.

In our study about 7% respondents do have no know the way of transmission on novel corona virus yet as well as the symptoms of the coronavirus. Thus, those who have no clue about the transmission of COVID 19, have chance to spread disease in the community. Thus, both the public organization, private sectors and concerned bodies should set strategies how to fill this knowledge gap. In addition, ever body should have played the great role in creating awareness about the about a coronavirus outbreak and to date, the virus is no drug or vaccine proven to treat or prevent COVID-19, in order to reduce the transmission of COVID-19 and to save many lives.

In general, as the data presented in the study discloses, the respondents expect continuous creation among the community and the zonal government and concerned bodies to pay attention on pandemic issues. On the other hand, most of the respondents indicated that, even the awareness was created among the community through different Media and different concerned bodies, only small portion of community have sufficient awareness. But large portion of community are not still totally pay attention to aware about COVID 19, crisis. Therefore, it seems reasonable to conclude that the community needs additional efforts from concerned body to address full awareness about COVID-19 pandemic.

Taking time to read, listen or watch informative media is an essential part of any professional's life and any bodies to prevent themselves from the deadly virus, but not everyone makes this part of their daily life. If leaders are not constantly eager to expand their knowledge, especially in their particular field, it would be so difficult to be successful in their leadership. Uncreated awareness can incur various challenges including poor choices, focus placed in the wrong areas increased costs, lost opportunities, decreased leverage, loss of efficiency, dissatisfaction. Ultimately poor planning impacts on the nature and quality of service provided because the resources are not available to meet existing needs or respond quickly to ever Changing needs. In a study conducted in the awareness of COVID-19 pandemic among interns by (Priya and Sherkhane 2020), tertiary care hospital in Dharwad district, Karnataka, were updating their knowledge on an everyday basis and knew the protocols and standard operating procedures framed by the state. This shows 26.8% of participants were still doubtful where the creation of awareness was created or not. This is the reason of being farness from electronics media and from the source of information as well as lack of skill to search relevant information from different social media through technology. In addition, certain communities are not careful and they did not follow the media effectively to aware about coronavirus and to prevent oneself as well as the community from such virus.

In this regard all of the informants (100%) have responded that the awareness was created among the community through different media and different place. The reason why crating awareness is an important for community is that, the proper aware of the crisis of COVID 19, protect the community from exposer and minimize the spreading virus throughout the community. It looks evident from the description of the data in the study shows that the awareness was created through different media or not.

According to study conducted by (Priya and Sherkhane, 2020), there is death in awareness of coronavirus pandemic which is masked by more of myths rather than authentic facts. In this study the results of the study are not depending facts and the results are seeming fabricated truth which does not follow scientific procedures to collect the data that comprises awareness. This in other way have no scientific base and the result may not be scientific. In our study participants were aware of the COVID 19, disease symptoms, the controlling process and the way of spreading in different degrees. This

finding was revealed by collecting data by following scientific procedures and hence, the result was scientific contrary to the above research finding. In other words, the majority of individuals are prohibited from engaging in commonplace actions such as hand shaking and kissing. In other words, people are forbidden from engaging in activities that promote the transmission of the coronavirus in general. As a result, communities are limited in their ability to shake hands during social events such as funerals, wedding, and so on.

This result on the other hand opposed with the most practical activity of most area i.e., many people were observed hand shaking and cares as well as unlimited mobility while data collection period against prevention of spread of covid-19 virus is to limit the community activities between people to prevent infection through the government's action rules (Wu Z, McGoogan JM., 2020), more over on the study done by Rahman & Sathi, 2020 in which 18-29 ages are more likely stay at home. The respondents in the study area about 149 (37.3%) reacted that strongly dis agree with respect to corona virus disease can be prevented by using alcohol and sanitizer. 146(36.5%) of the people in the study area do not relate Covid-19 viral disease with belief, whereas 85(21.3%) strongly agree on covid-19 disease is related with belief of the society in the area. Respondents in the study area replied that about 232 (58%) responded that using mask is not effective preventive measure of the covid-19 in the area. This may be due to due some people perceive that mask by itself cases respiratory health problem or personal belief that COVID-19 is devein curse and that cannot stop by wearing of face mask. Majority of the respondents 196(49%) responded it is possible to prevent covid-19 disease by using or practicing preventive guideline set by health organization.

V. CONCLUSION

The findings in this study revealed that community's awareness is a high on each specific aspect of COVID-19 among the study participants. However, only some member of community had adequate awareness about the COVID-19 disease. Community only acquired sufficient information about the preventing and controlling the spread of COVID-19 virus, its symptoms, way transmission and available treatments. Ethiopia's MOH's efforts increases public awareness of community about the novel virus by sending mobile messages and using different means of communication, including television and social media.

Communities are not still totally pay attention to aware about COVID 19, crisis. Therefore, it seems reasonable to conclude that the community needs additional efforts from concerned body to address full awareness about COVID-19 pandemic. Related to this taking time to read, listen or watch informative media is an essential part of any professional's life and any bodies to prevent themselves from the deadly virus, but not everyone makes this part of their daily life.

In this regard all of the informants (100%) have responded that the awareness was created among the community through different media and different place. The reason why creating awareness is an important for community is that, the proper aware on the crisis of COVID 19, protect the community from exposer and minimize the spreading virus throughout the community.

The reason why creating awareness is an important for community is that, the proper aware of the crisis of COVID 19, protect the community from exposer and minimize the spreading virus throughout the community. Communities have knowledge, awareness and confidence that Corona virus disease transmit from victim to healthy one during sneezing and coughing as they are aware from different ways.

The findings in this study revealed that community's awareness is a high on each specific aspect of COVID-19 among the study participants. However, only some member of community had adequate awareness about the COVID-19 disease. Community only acquired sufficient information about the preventing and controlling the spread of COVID-19 virus, its symptoms, way transmission and available treatments. Ethiopia's MOH's efforts increases public awareness of community about the novel virus by sending mobile messages and using different means of communication, including television and social media.

Community are not still totally pay attention to aware about COVID 19, crisis. Therefore, it seems reasonable to conclude that the community needs additional efforts from concerned body to address full awareness about COVID-19 pandemic. Related to this taking time to read, listen or watch informative media is an essential part of any professional's life and any bodies to prevent themselves from the deadly virus, but not everyone makes this part of their daily life.

In this regard all of the informants (100%) have responded that the awareness was created among the community through different media and different place. The reason why creating awareness is an important for community is that, the proper aware on the crisis of COVID 19, protect the community from exposer and minimize the spreading virus throughout the community.

The reason why creating awareness is an important for community is that, the proper aware of the crisis of COVID 19, protect the community from exposer and minimize the spreading virus throughout the community. Communities have knowledge and awareness and confidence that Corona virus disease transmit from victim to healthy one during sneezing and coughing as they are aware from different ways. And hence, the finding revealed that According to finding in this research communities wash their hand in order to prevent covid-19 better than the previous.

RECOMMENDATIONS

Some of the participants had no adequate awareness and preventive practices about the COVID-19 disease to protect one self, family as well as community and the world. Hence, Kaffa zone government should create adequate awareness as

well as to practice how to prevent oneself from COVID-19 by making dual cooperation with Kaffa zone health organizations. In addition, community only acquired sufficient information about how to preventing and control the spread of COVID-19 virus, its symptoms, way transmission and available treatments. What we suggest here is that approximately communities in Ethiopia settled in rural settlement where public infrastructures are less and even, we can say absent; among this electricity is the one by which peoples consume power for the sake of television watching. In spite of this, Kaffa zone as well as stakeholders should have to provide ample information for all communities including urban and rural societies to aware about COVID-19 and practice how to prevent.

All concerned bodies should have to develop the attention of the communities to keep all preventive practices that WHO have declared to protect from COVID-19.

Some peoples perceive that mask by itself causes respiratory health problem or personal belief that COVID-19 is devein curse and that cannot stop by wearing of face mask. Related to this Kaffa zone health organization and religious leaders should have to work cooperatively that using face mask can prevent us from COVID-19 by against this disease is not devein curse.

ACKNOWLEDGMENT

We would like to thank Bonga University because of funding to accomplish our study. Our deep gratitude also goes to our study subjects who were volunteered and took their time to give us all the relevant information for the study.

Competing interest

The authors have declared that no competing interests.

REFERENCES

- [1]. Adhikari SP., Meng S, Wu YJ, Mao YP, Ye RX, Wang QZ, Sun C, Sylvia S, Rozelle S and Raat H (2020). Epidemiology, causes, clinical manifestation and diagnosis, prevention and control of coronavirus disease (COVID-19) during the early outbreak period: a scoping review. *Infectious diseases of poverty*, 9(1): 1-12.
- [2]. Annune, A. E. (2020) 'Digital Commons @ University of Nebraska - Lincoln Sensitization and Awareness Creation as Tools for Curbing Perceived Effects of COVID-19 Pandemic on University Library Users in Nigeria', (September).
- [3]. Bhagavathula AS, Aldhaleei WA, Rahmani J, Mahabadi MA, Bandari DK. Novel Coronavirus (COVID-19) Knowledge and Perceptions: A Survey on Healthcare workers. *MedRxiv* 2020:2020.03.09.20033381. <https://doi.org/10.1101/2020.03.09.20033381>
- [4]. CDC:(2020). Interim Infection Prevention and Control Recommendations for Patients with Suspected or Confirmed Coronavirus Disease (COVID-19) in Healthcare Settings.

- [5]. Guo YR, Cao QD, Hong ZS, Tan YY, Chen SD, Jin HJ, Tan KS, Wang DY and Yan Y (2020). doi:10.1056/NEJMoa2001017 [[PMC](#) [free article](#)] [[PubMed](#)] [[Cross Ref](#)] [[Google Scholar](#)]
- [6]. The origin, transmission and clinical therapies on coronavirus disease 2019 (COVID-19) outbreak—an update on the status. *Military Medical Research* 7(1): 1-10.
- [7]. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; 395:497–506. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5).
- [8]. Hussain A, Garima T, Singh B, Ram R, Tripti R. Knowledge, attitudes, and practices towards COVID-19 among Nepalese residents: a quick online cross-sectional survey. *Asian J Med Sci.* 2020;11 (3):6–11. doi:10.3126/ajms.v11i3.2848519.
- [9]. Ioannidis JPA. Coronavirus disease 2019: the harms of exaggerated information and non-evidence-based measures. *Eur J Clin Invest.* 2020; e13223.
- [10]. Lu R, Zhao X, Li J, Niu P, Yang B, Wu H, et al. Genomic characterization and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet* 2020; 395:565–74. [https://doi.org/10.1016/S0140-6736\(20\)30251-8](https://doi.org/10.1016/S0140-6736(20)30251-8)
- [11]. Mantwill S, Monestel-Umanˆa S, Schulz PJ. The relationship between health literacy and health disparities: a systematic review. *PLoSOne.*2015; journal.pone.0145455.
- [12]. Nkengasong JN, Mankoula W. Looming threat of COVID-19 infection in Africa: act collectively, and fast. *Lancet.* 2020;395(10227):841–842. doi:10.1016/S0140-6736(20)30464-5 [[PMC](#) [free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)].
- [13]. Peng Y, Chenchen P, Zheng Y, et al. *Knowledge, Attitude and Practice Associated with COVID-19 Among University Students: A Cross-Sectional Survey in China.* 2020.
- [14]. Priya, P. and Sherkhane, M. S. (2020) ‘Awareness of coronavirus disease (COVID-19) pandemic among interns of a tertiary care hospital’, 9(6), pp. 375–381. doi: 10.5455/ijmsph.2020.06084222022062020.
- [15]. Voo T, Capps B. Influenza pandemic and the duties of healthcare professionals. *Singapore Med J.* 2010;51(4):275–281. [[PubMed](#)] [[Google Scholar](#)].
- [16]. Wolf MS, Gazmararian JA, Baker DW. Health literacy and functional health status among older adults. *Arch Intern Med.* 2005;165: 1946-52.
- [17]. Worldometer. COVID-19 coronavirus outbreak. Accessed at [www. Worldometer. info/coronavirus](http://www.Worldometer.info/coronavirus) on 24 March 2020.
- [18]. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *Jama.* 2020 Feb 24. <https://doi.org/10.1001/jama.2020.2648> PMID: 32091533.
- [19]. Zelalem T, Malede Y, Zenebe N, Akeberegn A, (2020), COVID-19-Related Knowledge, Attitude and Practice Among Hospital and Community, Dove Press journal:
- [20]. Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med.* 2020;382(8):727–733.